

Math Assignment Experts is a leading provider of online Math help. Our experts have prepared sample assignments to demonstrate the quality of solution we provide. If you are looking for mathematics help then share your requirements at info@mathassignmentexperts.com

Limits and Continuity

1 A tank contains 10 liters of pure water. Salt water containing 20 grams of salt per liter is pumped into the tank at 2 liters per minute.

1. Express the salt concentration $C(t)$ after t minutes (in g/L).
2. What is the long-term concentration of salt, i.e., $\lim_{t \rightarrow \infty} C(t)$?

Solution:

1. The concentration is, in units of g/L,

$$C(t) = \frac{\text{total salt}}{\text{total volume}} = \frac{20 \cdot 2 \cdot t}{10 + 2 \cdot t} = \boxed{\frac{20t}{5 + t}}$$

2. The long-term concentration is, in units of g/L,

$$\lim_{t \rightarrow \infty} \frac{20t}{5 + t} = \lim_{t \rightarrow +\infty} \frac{20t}{5 + t} \cdot \frac{1/t}{1/t} = \lim_{t \rightarrow +\infty} \frac{20}{5/t + 1} = \boxed{20}$$

2 Find the values of a and b that make $f(x)$ continuous for all real x .

$$f(x) = \begin{cases} be^x + a + 1, & x \leq 0 \\ ax^2 + b(x + 3), & 0 < x \leq 1 \\ a \cos(\pi x) + 7bx, & x > 1 \end{cases}$$

Solution: We note that the functions are continuous on their domains, so we check that the left- and right-hand limits agree at the boundary x -values. At $x = 0$,

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} be^x + a + 1 = b + a + 1,$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} ax^2 + b(x + 3) = 3b,$$

so $b + a + 1 = 3b$, and $a = 2b - 1$. Next, at $x = 1$,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} ax^2 + b(x + 3) = a + 4b,$$


$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} a \cos(\pi x) + 7bx = -a + 7b,$$

so $a + 4b = -a + 7b$, and $2a = 3b$. Solving this linear system in a and b yields $b = 2$ and $a = 3$ as the only solution.

3 Sketch the graph of a function f with the following properties:

- $\lim_{x \rightarrow 1} f(x) = 2$, but $f(1) = 1$
- $\lim_{x \rightarrow 3} f(x) = +\infty$
- $\lim_{x \rightarrow 2^+} f(x) = -1$, $\lim_{x \rightarrow 2^-} f(x) = 3$
- $\lim_{x \rightarrow +\infty} f(x) = -2$
- $\lim_{x \rightarrow -\infty} f(x) = -\infty$

Solution: Answers may vary, but here is a representative solution:



4 Show that the equation $\sqrt{x-5} = \frac{1}{x+3}$ has at least one real solution.

Solution: Let $f(x) = \sqrt{x-5} - \frac{1}{x+3}$, so that $f(x) = 0$ if and only if x is a solution to the equation. Then f is defined and continuous for all $x \geq 5$. Evaluating f at 5 and at 6, we see that

$$f(5) = \sqrt{5-5} - \frac{1}{5+3} = -\frac{1}{8} < 0 \quad \text{and} \quad f(6) = \sqrt{6-5} - \frac{1}{6+3} = \frac{8}{9} > 0.$$

By the Intermediate Value Theorem, there is some c in the interval $(5, 6)$ so that $f(c) = 0$, so f has at least one root.

(In fact, it is possible to reduce this equation to the cubic polynomial equation $(x-5)(x+3)^2 - 1 = 0$, and it is unpleasant but not impossible to find its roots exactly; the only valid root of the original equation is

$$c = -\frac{1}{3} + \frac{1}{3} \sqrt[3]{\frac{1051 - 15\sqrt{249}}{2}} + \frac{1}{3} \sqrt[3]{\frac{1051 + 15\sqrt{249}}{2}} \approx 5.01556 \dots .)$$

5 Consider the rational function

$$f(x) = \frac{x^5 - x^4 - 2x^3}{x^4 - 3x^3 - x^2 + 3x}$$

- For what values of a does f have a removable discontinuity at a ? What is $\lim_{x \rightarrow a} f(x)$ at those a ?
- For what values of a does f have an infinite discontinuity at a ?
- What is $\lim_{x \rightarrow +\infty} f(x)$?

(Hint: Factor the numerator and the denominator.)

Solution: We factor the numerator and denominator of f to obtain

$$f(x) = \frac{x^3(x+1)(x-2)}{x(x+1)(x-1)(x-3)}.$$

Away from $x = 0$ and $x = -1$, $f(x)$ simplifies to

$$\frac{x^2(x-2)}{(x-1)(x-3)},$$

which is defined and continuous at these x -values. Thus, f has removable discontinuities at $x = 0$ and at $x = -1$. From this form of f , we also compute that

$$\begin{aligned} \lim_{x \rightarrow 0} f(x) &= \lim_{x \rightarrow 0} \frac{x^2(x-2)}{(x-1)(x-3)} = \frac{0^2(-2)}{(-1)(-3)} = 0 \quad \text{and} \\ \lim_{x \rightarrow -1} f(x) &= \lim_{x \rightarrow -1} \frac{x^2(x-2)}{(x-1)(x-3)} = \frac{(-1)^2(-3)}{(-2)(-4)} = -\frac{3}{8}. \end{aligned}$$

At $x = 1$ and at $x = 3$, however, the numerator of the original function is not 0, so f has infinite singularities (and vertical asymptotes) at these x -values.

Finally, we observe that

$$\lim_{x \rightarrow +\infty} \frac{x^5 - x^4 - 2x^3}{x^4 - 3x^3 - x^2 + 3x} = \lim_{x \rightarrow +\infty} \frac{x - 1 - \frac{2}{x}}{1 - \frac{3}{x} - \frac{1}{x^2} + \frac{3}{x^3}} = +\infty.$$

6 Find the value of a such that

$$\lim_{x \rightarrow -1} \frac{2x^2 - ax - 14}{x^2 - 2x - 3}$$

exists. What is the value of the limit?

Solution: We observe that $\lim_{x \rightarrow -1} x^2 - 2x - 3 = 1 + 2 - 3 = 0$, so in order for this limit to exist, we need the limit of the numerator as $x \rightarrow -1$ to be 0 as well. Since

$$\lim_{x \rightarrow -1} 2x^2 - ax - 14 = 2 + a - 14 = a - 12,$$

$a - 12 = 0$, and $a = 12$. Then, away from $x = -1$,

$$\frac{2x^2 - 12x - 14}{x^2 - 2x - 3} = \frac{2(x - 7)(x + 1)}{(x - 3)(x + 1)} = \frac{2(x - 7)}{(x - 3)}.$$

As $x \rightarrow -1$, we see by evaluation that the limit is $\frac{2(-8)}{(-4)} = 4$.